arXiv:1304.7791v1 [astro-ph.CO] 29 Apr 2013

Can all cosmological observations be accurately interpreted with a unique geometry?

Pierre Fleury[1,3],* Hélène Dupuy[1,2,3],† and Jean-Philippe Uzan[1,3]‡

[1] *Institut d'Astrophysique de Paris, UMR-7095 du CNRS,*
Université Pierre et Marie Curie, 98 bis bd Arago, 75014 Paris, France.
[2] *Institut de Physique Théorique, CEA, IPhT, URA 2306 CNRS, F-91191 Gif-sur-Yvette, France.*
[3] *Sorbonne Universités, Institut Lagrange de Paris, 98 bis bd Arago, 75014 Paris, France.*

(Dated: May 1, 2013)

The recent analysis of the *Planck* results reveals a tension between the best fits for $(\Omega_{\rm m0}, H_0)$ derived from the cosmic microwave background or baryonic acoustic oscillations on the one hand, and the Hubble diagram on the other hand. These observations probe the universe on very different scales since they involve light beams of very different angular sizes, hence the tension between them may indicate that they should not be interpreted the same way. More precisely, this letter questions the accuracy of using only the (perturbed) Friedmann-Lemaître geometry to interpret all the cosmological observations, regardless of their angular or spatial resolution. We show that using an inhomogeneous "Swiss-cheese" model to interpret the Hubble diagram allows to reconcile it with the *Planck* results. Such an approach does not require to invoke new physics nor to violate the Copernican principle.

PACS numbers: 98.80.-k, 04.20.-q, 42.15.-i.

The standard interpretation of cosmological data relies on the description of the universe by a spatially homogeneous and isotropic spacetime with a Friedmann-Lemaître (FL) geometry, allowing for perturbations [1]. The emergence of a dark sector, including dark matter and dark energy, emphasizes the need for extra-degrees of freedom, either physical (new fundamental fields or interactions) or geometrical (e.g. a cosmological solution with lower symmetry). This has driven a large activity to test the hypotheses [2] of the cosmological model, such as general relativity or the Copernican principle.

The recent *Planck* data were analyzed in such a framework [3] in which the cosmic microwave background (CMB) anisotropies are treated as perturbations around a FL universe, with most of the analysis performed at linear order. Non-linear effects remain small [4] and below the constraints on non-Gaussianity derived by *Planck* [5]. The results nicely confirm the standard cosmological model of a spatially Euclidean FL universe with a cosmological constant, dark matter and initial perturbations compatible with the predictions of inflation.

Among the constraints derived from the CMB, the Hubble parameter H_0 and the matter density parameter $\Omega_{\rm m0}$ are mostly constrained through the combination $\Omega_{\rm m0}h^3$, where $H_0 = 100h$ km/s/Mpc. It is set by the acoustic scale $\theta_* = r_{\rm s}/D_{\rm A}$, defined as the ratio between the sound horizon and the angular distance at the time of last-scattering. The measurement of seven acoustic peaks allows to determine θ_* with a precision better than 0.1%. The constraints on the plane $(\Omega_{\rm m0}, H_0)$ are presented in Fig. 3 of Ref. [3] and clearly show this degeneracy. The marginalized constraints on the two parameters were then derived [3] to be

$$H_0 = (67.3\pm1.2)\ \text{km/s/Mpc}, \quad \Omega_{\rm m0} = 0.315\pm0.017 \quad (1)$$

at a 68% confidence level. It was pointed out (see § 5.2–5.4 of Ref. [3]) that the values of H_0 and $\Omega_{\rm m0}$ are respectively low and high compared with their values inferred from the Hubble diagram. Such a trend was already indicated by WMAP-9 [6] which concluded $H_0 = (70 \pm 2.2)$ km/s/Mpc.

Regarding the Hubble constant, two astrophysical measurements are in remarkable agreement. First, the estimation based on the distance ladder calibrated by three different techniques (masers, Milky way cepheids and Large Magelanic cloud cepheids) gives [7] $H_0 = (74.3 \pm 1.5 \pm 2.1)$ km/s/Mpc, respectively with statistical and systematic errors. This improves the earlier constraint obtained by the HST Key program [8], $H_0 = (72 \pm 8)$ km/s/Mpc. Second, the Hubble diagram of type Ia supernovae (SNe Ia) calibrated with the HST observations of cepheids leads [9] to $H_0 = (73.8\pm2.4)$ km/s/Mpc. Other determinations of the Hubble constant, e.g. from VLBI observations [10] or from the combination of Sunyaev-Zel'dovich effect and X-ray observations [11], have larger error bars and are compatible with both the CMB and distance measurements.

Besides, the analysis of the Hubble diagram of SNe Ia leads to a lower value of $\Omega_{\rm m0}$ – e.g. 0.222 ± 0.034 with the SNLS 3 dataset [12] – compared to the constraint (1) by *Planck*. As concluded in Ref. [3], there is no direct inconsistency, and it was pointed out that there could be residual systematics not properly accounted for in the SNe data. Still, it was stated that "the tension between CMB-based estimates and the astrophysical measurements of H_0 is intriguing and merits further discussion".

Interestingly, the CMB constraints on $(\Omega_{\rm m0}, H_0)$ are in excellent agreement with baryon acoustic oscillation

---

*Electronic address: fleury@iap.fr
†Electronic address: helene.dupuy@cea.fr
‡Electronic address: uzan@iap.fr

(BAO) measurements [13], that allow to determine the angular distance up to redshifts of order 0.7. The common point between the CMB and BAO measurements is that they involve light beams much larger than those involved in astronomical observations. Indeed, a pixel of *Planck*'s high-resolution CMB maps corresponds to 5 arcmin [14], while the typical angular size of a SN is 10^{-7} arcsec. This means that *the two kinds of observations probe the universe at very different scales.* Moreover, for both the CMB and BAO measurements the crucial information is encoded in correlations, while SNe observations rely on "1-point measurements" (we are interested in the luminosity/redshift of each SN, not in the correlations between several SNe). Because of such distinctions one can expect from the two classes of cosmological observations to be affected differently by the inhomogeneity of the universe, through gravitationnal lensing.

The effect of lensing on CMB measurements is essentially due to the large-scale structure, and it can be taken into account in the framework of cosmological perturbation theory at linear order [15] (see however Ref. [16] for a discussion about the impact of strong inhomogeneities). We refer to Ref. [17] for a description of the lensing effects on BAO measurements. Regarding the Hubble diagram, the influence of lensing has also been widely investigated [18]. The propagation of light in an inhomogeneous universe gives rise to both distortion and magnification. In principle, most images are demagnified because their lines of sight probe underdense regions, while some are amplified due to strong lensing. It shall thus induce a dispersion of the luminosities of the sources, that is, an extra-scatter in the Hubble diagram [19]. Its amplitude can be determined from the perturbation theory [20] and substracted [21]. However, a considerable fraction of the lensing effects arises from subarcminute scales, which are not probed by shear maps smoothed on arcminute scales [22].

The tension on $(\Omega_{\mathrm{m}0}, H_0)$ may indicate that, given the accuracy of the observations achieved today, the use of a (perturbed) FL geometry to interpret the astrophysical data is no longer adapted. More precisely, the question that we want to raise is *whether the use of a unique space-time geometry is relevant for interpreting all the cosmological observations, regardless of their angular or spatial resolution and of their location (redshift).* Indeed, each observation is expected to probe the universe smoothed on a typical scale related to its resolution, and this can lead to fundamentally different geometrical situations. In a universe with a discrete distribution of matter, the Riemann curvature experienced by a beam of test particles or photons is dominated by the Weyl tensor. Conversely, in a (statistically spatially isotropic) universe smoothed on large scales it is dominated by the Ricci tensor. Both situations correspond to distinct optical properties [23].

In the framework of geometric optics, a light beam is described by a bundle of null geodesics. All the information about the size and the shape of a beam can be encoded in a 2×2 matrix $\mathcal{D}^A{}_B$ called the Jacobi map (see Ref. [24] for futher details). In particular, the angular and luminosity distances read respectively

$$D_{\mathrm{A}} = \sqrt{|\det \mathcal{D}^A{}_B|}, \qquad D_{\mathrm{L}} = (1+z)^2 D_{\mathrm{A}}. \tag{2}$$

where z denotes the redshift. The evolution of the Jacobi map with light propagation is governed by the Sachs equation [25, 26]

$$\frac{\mathrm{d}^2}{\mathrm{d}v^2}\,\mathcal{D}^A{}_B = \mathcal{R}^A{}_C\,\mathcal{D}^C{}_B, \tag{3}$$

where v is an affine parameter along the geodesic bundle. The term $\mathcal{R}_{AB}$ which controls the evolution of $\mathcal{D}^A{}_B$ is a projection of the Riemann tensor called the optical tidal matrix. It is defined by $\mathcal{R}_{AB} \equiv R_{\mu\nu\alpha\beta} k^\nu k^\alpha s^\mu_A s^\beta_B$, where k^μ is the wave vector of an arbitrary ray, and the Sachs basis $\{s^\mu_A\}_{A=1,2}$ spans a screen on which the observer projects the light beam. Because the Riemann tensor can be split into a Ricci part and a Weyl part, the optical tidal matrix can also be decomposed as

$$(\mathcal{R}_{AB}) = \underbrace{\begin{pmatrix} \Phi_{00} & 0 \\ 0 & \Phi_{00} \end{pmatrix}}_{\text{Ricci lensing}} + \underbrace{\begin{pmatrix} -\mathrm{Re}\,\Psi_0 & \mathrm{Im}\,\Psi_0 \\ \mathrm{Im}\,\Psi_0 & \mathrm{Re}\,\Psi_0 \end{pmatrix}}_{\text{Weyl lensing}} \tag{4}$$

with $\Phi_{00} \equiv -(1/2) R_{\mu\nu} k^\mu k^\nu$ and $\Psi_0 \equiv -(1/2) C_{\mu\nu\alpha\beta}(s^\mu_1 - \mathrm{i} s^\mu_2) k^\nu (s^\beta_1 - \mathrm{i} s^\beta_2) k^\beta$. It clearly appears in Eq. (4) that the Ricci term tends to isotropically focus the light beam, while the Weyl term tends to shear and rotate it. The behaviour of a light beam is thus different whether it experiences Ricci-dominated lensing (large beams, e.g. CMB measurements) or Weyl-dominated lensing (narrow beams, e.g. SNe observations). It has been demonstrated [27] that for the lensing caused by a distribution of pointlike particles, the Ricci-lensing behaviour is indeed recovered when the size of the beam increases, as the light beam encloses more and more particles.

This Ricci-Weyl problem can be addressed with different methods. One possibility, a representative of which is the Dyer-Roeder approximation [28], is to construct a general distance-redshift relation which would take into account the effect of inhomogeneities in some average way. However, such approaches are in general difficult to control [18] because they rely on approximations whose domain of applicability is unknown. An alternative possibility consists in constructing inhomogeneous cosmological models, with a discrete distribution of matter, and studying the impact on light propagation. Several models exist in the litterature: the Schwarzschild-cell method [29] or the lattice universe [30] which are both approximate solutions of the Einstein's equations; and the Swiss-cheese models [31] which are constructed by matching together patches of exact solutions of the Einstein's equations. This last approach is the one that we shall follow in the present letter.

Consider a Swiss-cheese model in which clumps of matter, each of them lying at the center of a spherical void, are embedded in a FL spacetime. The interior region of a void is thus described by the Kottler geometry (i.e. Schwarzschild with a cosmological constant), while the exterior geometry is the FL one. By construction, such inhomogeneities do not modify the expansion dynamics of the embedding FL universe, thus avoiding any discussion regarding backreaction. The resulting spacetime is well-defined, because the Darmois-Israel junction conditions are satisfied on the boundary of every void. Compared to a striclty homogeneous universe, a Swiss-cheese model is therefore characterized by two additional parameters: the size of the voids (or equivalently the mass M of their central bodies), and the volumic fraction of the remaining FL regions which encodes the smoothness of the distribution of matter. It is naturally quantified by

$$f \equiv \lim_{V\to\infty} \frac{V_{\rm FL}}{V}, \tag{5}$$

called *smoothness parameter*, where $V_{\rm FL}$ is the volume occupied by the FL region within a volume V of the Swiss-cheese. With the definition (5), $f = 1$ corresponds to a model with no hole – i.e. a FL universe – while $f = 0$ corresponds to the case where matter is exclusively under the form of clumps.

The propagation of light in this model has been comprehensively investigated in Ref. [24] as well as the effects on the Hubble diagram, generalizing earlier works [32]. Note that the case of voids with a Lemaître-Tolman-Bondi geometry has also been considered in the litterature, but they were shown to have only a marginal effect on the Hubble diagram [33] (and it does not question the fluid limit). Of course Swiss-cheese models cannot be considered as realistic cosmological models, but neither do the exact FL geometry used to interpret the Hubble diagram. Both spacetimes describe a spatially *statistically* homogeneous and isotropic universe, and the former allows additionally to investigate the effect of a discrete distribution of matter. Since the FL universe is a particular Swiss-cheese model, this family of spacetimes therefore allows to estimate how good the hypothesis of strict spatial homogeneity (with a continuous matter distribution at *all* scales) is.

In our previous analysis [24], we integrated the Sachs equation (3) and simulated Hubble diagrams for a series of Swiss-cheese models with various cosmological parameters $(\Omega_{\rm m0}, \Omega_{\Lambda 0})$ characterizing the FL region, and for different smoothness parameters f. This allowed us to determine the typical error on the parameters induced by wrongly assuming that light propagates in a FL spacetime, when the observations are interpreted. The effect of inhomogeneities is of order of a few percents, which can be significant given the precision on the cosmological parameters reached today. Such an order of magnitude is comparable with other estimates in similar contexts [34].

The synthetic Hubble diagrams enabled us to infer a phenomenological expression for the luminosity distance $D_{\rm L}(z; \Omega_{\rm m0}, \Omega_{\Lambda 0}, H_0, f)$ depending on the smoothness parameter f. This expression was then used to fit the Hubble diagram constructed from the SNLS 3 catalog [12]. Fig. 25 of Ref. [24] shows that f influences the result of the best-fit on $\Omega_{\rm m0}$ that can shift from 0.22 for $f = 1$ (in agreement with the standard FL analysis performed in Ref. [12]) to 0.3 for $f = 0$.

We present here the constraints in the plane $(H_0, \Omega_{\rm m0})$ for different values of f (assumed to be constant). The result is depicted on Fig. 1 which clearly shows that the agreement between the CMB and the Hubble diagram is improved for small values of f. We also note that H_0 decreases with f, but too slightly to completely reconcile all the observations.



FIG. 1: Comparison of the constraints obtained by *Planck* on $(\Omega_{\rm m0}, H_0)$ [3] and from the analysis of the Hubble diagram constructed from the SNLS 3 catalog. The different contour plots correspond to different smoothness parameters f. For $f = 1$, the geometry used to fit the data is Friedmannian, as assumed in standard analysis.

In principle, the smoothness parameter f should be related to the growth of structures, so that we expect it to be an increasing function of z. A generic form

$$f(z) = 1 + \frac{f_0 - 1}{1 + z/z_*}, \tag{6}$$

which interpolates between f_0 today and $f = 1$ at high redshift, can be used to take this effect into account. In this case, the impact of the inhomogeneities on the Hubble diagram is weaker. A redshift-dependent smoothness therefore reduces the agreement between the Hubble diagram and the CMB compared to the case $f = f_0$, or alternatively requires a lower value for f_0.

Our analysis, though relying on a particular class of models, indicates that the FL geometry is probably too simplistic to describe the universe for certain types of observations, given the accuracy reached today. In the end, a single metric may not be sufficient to describe all the cosmological observations, just as *Liliputians* and *Brobdingnag*'s giants [35] cannot use a map with the same resolution to travel. A better cosmological model probably requires an atlas of maps with various smoothing scales, determined by the observations at hand.

Other observations, such as lensing [36], may help to characterize the distribution and geometry of voids [37], in order to construct a better geometrical model. For the first time, the standard FL background geometry may be showing its limits to interpret the cosmological data with the accuracy they require.

Acknowledgements. We thank Francis Bernardeau, Yannick Mellier, Alice Pisani, Cyril Pitrou, Joe Silk and Benjamin Wandelt for discussions. This work was supported by French state funds managed by the ANR within the Investissements d'Avenir programme under reference ANR–11–IDEX–0004–02, the Programe National Cosmologie et Galaxies, and the ANR THALES (ANR-10-BLAN-0507-01-02).

[1] P. Peter and J.-P. Uzan, *Primordial cosmology*, (Oxford Univ. Press, England, 2009).
[2] J.-P. Uzan, Gen. Rel. Grav. **39**, 307 (2007); J.-P. Uzan, `arXiv:0912.5452`; J.-P. Uzan, Living Rev. Rel. **14**, 2 (2011).
[3] Planck Collaboration, paper XVI, `arXiv:1303.5076`
[4] C. Pitrou, J.-P. Uzan, and F. Bernardeau, Phys. Rev. D **78**, 063526 (2008); C. Pitrou, J.-P. Uzan, and F. Bernardeau, JCAP **1007**, 003 (2010); Z. Huang, and F. Vernizzi, Phys. Rev. Lett. **110**, 101303 (2013).
[5] Planck Collaboration, paper XXIV, `arXiv:1303.5084`
[6] G. Hinshaw, *et al.*, `arXiv:1212.5226`
[7] W.L. Freedman, *et al.*., Astrophys. J. **758**, 24 (2012).
[8] W.L. Freedman, *et al.*., Astrophys. J. **553**, 47 (2001).
[9] A.G. Riess, *et al.*, Astrophys. J. **730**, 119 (2011).
[10] M.J. Reid, *et al.*, `arXiv:1207.7292`
[11] M. Bonamente, *et al.*, Astrophys. J. **647** 25 (2006).
[12] A. Conley, *et al.*, Astrophys. J. Suppl. **192**, 1 (2011).
[13] S. Cole, *et al.*, Month. Not. R. Astron. Soc. **362**, 505 (2005); D.J. Eisenstein, *et al.*, Astrophys. J. **633**, 560 (2005); W.J. Percival, *et al.*, Month. Not. R. Astron. Soc. **401**, 2148 (2011); N. Padmanabhan, *et al.*, Month. Not. R. Astron. Soc. **427**, 2132 (2012).
[14] Planck Collaboration, paper I, `arXiv:1303.5062`
[15] Planck Collaboration, paper XVII, `arXiv:1303.5077`
[16] K. Bolejko, JCAP **02**, 025 (2011).
[17] A. Vallinotto, S. Dodelson, C. Schimd and J.-P. Uzan, Phys. Rev. D **75**, 103509 (2007).
[18] C. Clarkson, *et al.*, Month. Not. R. Astron. Soc. **426**, 1121 (2012).
[19] R. Kantowski, T. Vaughan and D. Branch, Astrophys. J. **447**, 35 (1995); J.A. Frieman, Comm. Astrophys. **18**, 323 (1998); Y. Wang, Astrophys. J. **531**, 676 (2000).
[20] C. Bonvin, R. Durrer and M. A. Gasparini, Phys. Rev. D **73**, 023523 (2006); N. Meures and M. Bruni, `arXiv:1107.4433`; P. Valageas, Astron. Astrophys. **354**, 767, (2000).
[21] A. Cooray, D. Holz and D. Huterer, Astrophys. J. **637**, L77 (2006); S. Dodelson and A. Vallinotto, Phys. Rev. D **74**, 063515 (2006); A. Cooray, D. Huterer and D. Holz, Phys. Rev. Lett. **96**, 021301 (2006); D. Sarkar, *et al.*, `arXiv:0710.4143`; A. R. Cooray, D. E. Holz and R. Caldwell, JCAP **1011**, 015 (2010); A. Vallinotto, S. Dodelson and P. Zhang, `arXiv:1009.5590`. B. Ménard *et al.*, Astron. Astrophys. **403**, 817 (2003).
[22] N. Dalal, *et al.*, Astrophys. J. **585**, L11 (2003).
[23] C.C. Dyer, and R.C. Roeder, Gen. Relat. Grav. **13**, 1157 (1981).
[24] P. Fleury, H. Dupuy, and J.-P. Uzan, `arXiv:1302.5308`.
[25] R. Sachs, Proc. Roy. Soc. Lond. A **264**, 309 (1961).
[26] P. Schneider, J. Ehlers, and E. E. Falco, *Gravitational Lenses* (Springer, 1992).
[27] A.O. Petters, B. Rider, and A.M. Teguia, J. Math. Phys. **50**, 122501 (2009).
[28] C. C. Dyer, and R. C. Roeder, Astrophy. J. **174**, L115 (1972); C. C. Dyer, and R. C. Roeder, Astrophy. J. **180**, L31 (1973).
[29] R.W. Lindquist, and J.A. Wheeler, Rev. Mod. Phys. **29**, 432 (1957); T. Clifton, and P.G. Ferreira, Phys. Rev. D **80**, 103503 (2009); T. Clifton, and P.G. Ferreira, JCAP **10**, 26 (2009),
[30] D. E. Holz and R. M. Wald, Phys. Rev. D **58**, 063501 (1998); J.-P. Bruneton, and J. Larena, `arXiv:1204.3433`; J.-P. Bruneton, and J. Larena, `arXiv:1208.1411`; K. Kainulainen and V. Marra, Phys. Rev. D **80**, 127301 (2009).
[31] A. Einstein, and E.G. Straus, Rev. Mod. Phys. **17**, 120 (1945); A. Einstein, and E.G. Straus, Rev. Mod. Phys. **18**, 148 (1945).
[32] R. Kantowski, Astrophys. J. **155**, 89 (1969); C. C. Dyer, and R. C. Roeder, Astrophys. J. **189**, 167 (1974).
[33] N. Brouzakis, N. Tetradis and E. Tzavara, JCAP **0702**, 013 (2007); V. Marra, *et al.*, Phys. Rev. D **76**, 123004 (2007); T. Biswas and A. Notari, JCAP **0806**, 021 (2008); R. A. Vanderveld, E. E. Flanagan and I. Wasserman, Phys. Rev. D **78**, 083511 (2008); N. Brouzakis, N. Tetradis and E. Tzavara, JCAP **0804**, 008 (2008); T. Clifton and J. Zuntz, `arXiv:0902.0726`; W. Valkenburg, JCAP **0906**, 010 (2009); S. J. Szybka, Phys. Rev. D **84**, 044011 (2011); E. E. Flanagan, *et al.*, Phys. Rev. D **85**, 023510 (2012).
[34] K. Bolejko, and P.G. Ferreira, JCAP **1205**, 003 (2012).
[35] J. Swift, *Travels into Several Remote Nations of the World, in Four Parts. By Lemuel Gulliver, First a Surgeon, and then a Captain of Several Ships.*
[36] K. Bolejko, *et al.*, Phys. Rev. Lett. **110** 021302 (2013).
[37] G. Lavaux, and B.D. Wandelt, Mon. Not. Roy. Astron. Soc.**1392**, 403 (2010).